

**Instructions to the Students**

- Write only question numbers clearly outside the margin (1, 2, 3.i, 5.b, 4.c.ii, etc.).
- Do not write questions or any titles. (For ex. - Do not write **II. Answer the following**).
- After every answer, give a one-line space.
- For Multiple choice Questions - Both Option and Answer should be written.
- Bullet points & Sub-points should be written inside the margin.
- Do not fold / staple the paper.

**Section A**

1. The fragrance of the aldehyde and ketone are used for perfume and similar uses depend on : [ 1 ]
  - a) on its reactivity with other functional groups.
  - b) size and solubility of the aldehyde and ketone molecule.
  - c) only solubility of aldehydes and ketones.
  - d) moisture of the air.
2. Which one is the correct increasing order of bond strength of C-X bond? [ 1 ]
  - a) C-I < C-Br < C-Cl < C-F
  - b) C-F < C-Cl < C-Br < C-I
  - c) C-Cl < C-F < C-I < C-Br
  - d) C-Br < C-I < C-F < C-Cl
3. The correct IUPAC name of  $[\text{Pt}(\text{NH}_3)_2\text{Cl}_2]^{2+}$  is [ 1 ]
  - a) Diamminedichloridoplatinum (II)
  - b) Diamminedichloridoplatinum (IV)
  - c) Diamminedichloridoplatinum (O)
  - d) Diamminedichloridoplatinate (IV)
4. Aniline does not undergo Friedel–Crafts alkylation or acylation because: [ 1 ]
  - a) The  $-\text{NH}_2$  group is strongly activating and causes polyalkylation.
  - b) The  $-\text{NH}_2$  group withdraws electrons and deactivates the ring.
  - c) The  $-\text{NH}_2$  group reacts with the catalyst ( $\text{AlCl}_3$ ), which deactivates the ring
  - d) The reaction mixture produces too much heat and decomposes.
5. On mixing 30 ml of acetone with 20 ml of chloroform, the total volume of solution is: [ 1 ]
  - a) equal to 10 ml
  - b) less than 50 ml
  - c) greater than 50 ml
  - d) equal to 50 ml
6. Which of the following molecules has a chiral centre correctly labelled with an asterisk(\*)? [ 1 ]
  - a)  $\text{CH}_3\text{C}^*\text{HBrCH}_3$
  - b)  $\text{CH}_3\text{C}^*\text{HClCH}_2\text{Br}$
  - c)  $\text{HOCH}_2\text{C}^*\text{H}(\text{OH})\text{CH}_2\text{OH}$
  - d)  $\text{CH}_3\text{C}^*\text{Br}_2\text{CH}_3$
7. Which of the following compound will not undergo azo coupling reaction with benzene diazonium chloride ? [ 1 ]
  - a) Phenol
  - b) Aniline
  - c) Nitrobenzene
  - d) Anisole
8. Which of the following statements is not true about glucose ? [ 1 ]
  - a) It is an aldohexose.
  - b) On heating with HI it forms n-hexane.
  - c) It exists in furanose form.
  - d) It does not give Schiff's test.



15. **Assertion (A):** All naturally occurring  $\alpha$ -amino acids except glycine are optically active. [ 1 ]  
**Reason (R) :** Most naturally occurring amino acids have L-configuration  
a) Both (A) and (R) are true and (R) is the correct explanation of (A)  
b) Both (A) and (R) are true but (R) is not the correct explanation of (A)  
c) (A) is correct but (R) is wrong  
d) (A) is wrong but (R) is correct

16. **Assertion(A) :** An electrochemical cell can be set-up only if the redox reaction is spontaneous. [ 1 ]  
**Reason(R) :** A reaction is spontaneous if the free energy change is negative.  
a) Both (A) and (R) are true and (R) is the correct explanation of (A)  
b) Both (A) and (R) are true but (R) is not the correct explanation of (A)  
c) (A) is correct but (R) is wrong  
d) (A) is wrong but (R) is correct


### Section B

- 17.I. (a) A and B liquids on mixing show rise in temperature. Which type of deviation from Raoult's law is there and why ? [ 2 ]  
(b) Why can azeotropic mixture not be separated by fractional distillation ?

(OR)

- 17.II. I. When 50 mL of phenol and 50 mL of aniline are mixed, predict whether the volume of the solution is equal to, greater than or less than 100 mL. Give reason to support your answer. [ 2 ]  
II. Ritesh suggested adding salt to the box containing ice. He said this would keep the cold drink bottles cold for a longer time. How will Ritesh justify his suggestion?

18. Consider the graph for the reaction [ 2 ]  
 $\text{H}_2 + \text{I}_2 \rightleftharpoons 2\text{HI}$



- I. Calculate enthalpy of reaction and activation energy for the backward reaction.  
II. How will the catalyst affect the rate of this reaction? Explain.
19. In a chemistry practical class, the teacher gave his students an amine 'X' having molecular formula  $\text{C}_2\text{H}_7\text{N}$ , and asked the students to identify the type of amine. One of the students, Neeta, observed that it reacts with  $\text{C}_6\text{H}_5\text{SO}_2\text{Cl}$ , to give a compound which dissolves in NaOH solution. Can you help Neeta to identify the compound 'X'? [ 2 ]
20. When a coordination compound  $\text{CrCl}_3 \cdot 6\text{H}_2\text{O}$  is mixed with  $\text{AgNO}_3$ , two moles of  $\text{AgCl}$  are precipitated per mole of the compound. What is the structural formula and IUPAC name of the compound? [ 2 ]

21. Explain the mechanism of acid catalysed hydration of an alkene to form corresponding alcohol. [ 2 ]

### Section C

22. A solution of glucose (molar mass  $180 \text{ g mol}^{-1}$ ) in water has a boiling point of  $100.20^\circ\text{C}$ . Calculate the freezing point of the same solution. Molal constants for water  $K_f$  and  $K_b$  are  $1.86 \text{ K kg mol}^{-1}$  and  $0.512 \text{ K kg mol}^{-1}$  respectively. [ 3 ]

23. Calculate  $E_{\text{cell}}$  of a galvanic cell in which the following reaction takes place at  $25^\circ\text{C}$  : [ 3 ]  
 $\text{Zn(s)} + \text{Pb}^{2+}(0.02 \text{ M}) \longrightarrow \text{Zn}^{2+}(0.1 \text{ M}) + \text{Pb(s)}$   
 [Given :  $E^\circ_{\text{Zn}^{2+}/\text{Zn}} = -0.76 \text{ V}$ ,  $E^\circ_{\text{Pb}^{2+}/\text{Pb}} = -0.13 \text{ V}$ ;  
 $\log 2 = 0.3010$ ,  $\log 4 = 0.6021$ ,  $\log 5 = 0.6990$ ].

24. Explain the following: [ 3 ]





I. Toluene on treatment with  $\text{Cl}_2$  in sunlight gives benzyl chloride whereas when treated with  $\text{Cl}_2$  in dark gives o-chlorobenzene and p-chlorobenzene.

II. Finkelstein reaction is carried out in the presence of dry acetone.

III. neo pentylchloride has lower boiling point than isopentylchloride.

25. (i) What is Lanthanoid contraction? [ 3 ]  
 (ii) In which way the electronic configuration of d block elements differs from representative elements?  
 (iii) Calculate the magnetic moment of  $\text{Mn}^{2+}$  in Bohr Magneton.

26. Predict the products of the following reactions.(any 3) [ 3 ]



27. How are the following conversions carried out? [ 3 ]  
 (i) Benzyl chloride to Benzyl alcohol  
 (ii) Ethyl magnesium chloride to Propan-1-ol  
 (iii) Methyl mag. bromide to 2-Methylpropan-2-ol.

28. (i) Name two elements of 3d series which show anomalous electronic configuration. [ 3 ]  
 (ii) Write the preparation of  $\text{KMnO}_4$  from Pyrolusite ore ( $\text{MnO}_2$ )

### Section D


29. Rahul set-up an experiment to find resistance of aqueous KCl solution for different concentrations at 298 K using a conductivity cell connected to a Wheatstone bridge. He fed the Wheatstone bridge with A.C. power in the audio frequency range 0 to 5000 cycles per second. Hence the resistance was calculated from null point, he also calculated the conductivity  $\kappa$  and molar conductivity  $\Lambda_m$  and recorded his readings in tabular form.

| S.No | Conc. (M) | $\kappa \text{ S cm}^{-1}$ | $\Lambda_m \text{ S cm}^2 \text{ mol}^{-1}$ |
|------|-----------|----------------------------|---|
| 1.   | 1.00      | $111.3 \times 10^{-3}$     | 111.3                                       |
| 2.   | 0.10      | $12.9 \times 10^{-3}$      | 129.0                                       |
| 3.   | 0.01      | $1.41 \times 10^{-3}$      | 141.0                                       |


- 29.I. Why does conductivity decrease with dilution? [ 1 ]
- 29.II. If  $\Lambda_m^0$  of KCl is  $150.0 \text{ S cm}^2 \text{ mol}^{-1}$ , calculate the degree of dissociation of 0.01 M KCl. [ 1 ]
- 29.III. If Rahul had used HCl instead to KCl then would you expect the  $\Lambda_m$  values to be more or less than those per KCl for a given concentration. Justify [ 2 ]
30. In spite of the predictions of stable noble gas compounds since at least 1902, unsuccessful attempts at their synthesis gave rise to the widely held opinion that noble gases are not only noble but also inert. It was not until 1962 that this dogma was shattered when Bartlett in Canada published the first stable noble gas compound  $\text{XePtF}_6$ . This discovery triggered a worldwide frenzy in this area, and within a short time span many new xenon, radon, and krypton compounds were prepared and characterized. The recent discoveries show the ability of xenon to act as a ligand. The discovery by Seppelt's group that more than one xenon atom can attach itself to a metal center which in the case of gold leads to surprisingly stable Au- Xe bonds. The bonding in  $[\text{AuXe}_4]^{2+}$  involves 4 Xe ligands attached by relatively strong bonds to a single Au(II) center in a square planar arrangement with a Xe-Au bond length of about 274 pm This discovery provides not only the first example of multiple xenon ligands but also represents the first strong metal - xenon bond.  
 (Source: Christie, K. O. (2001). A renaissance in noble gas chemistry. Angewandte Chemie International Edition, 40(8), 1419-1421.)
- 30.I. What ligand is Xe in the given complex ? [ 1 ]
- 30.II. What is the hybridization shown by Au in  $[\text{AuXe}_4]^{2+}$  ? [ 1 ]
- 30.III. What is the difference between a geometrical isomer and an optical isomer in coordination compounds? [ 2 ]

## Section E

- 31.I. (i) Even though the phenoxide ion has more resonance structures than the carboxylate ion, carboxylic acids are stronger acids than phenol. Explain why ? [ 5 ]  
(ii) During the preparation of esters from a carboxylic acid and an alcohol in the presence of an acid catalyst, the water or the ester should be removed as soon as it is formed. Explain why ?  
(iii) Give one chemical test to distinguish Phenol and Acetic acid.  
(iv) Complete each synthesis by giving missing starting material, reagent or products



(a)



(b)

(OR)

- 31.II. (i)  $(\text{CH}_3)_3\text{C}-\text{O}-\text{CH}_3$  on reaction with  $\text{HI}$  gives  $(\text{CH}_3)_3\text{C}-\text{I}$  and  $\text{CH}_3-\text{OH}$  as the main products and not  $(\text{CH}_3)_3\text{C}-\text{OH}$  and  $\text{CH}_3-\text{I}$ ? [ 5 ]  
(ii) Give one chemical test to distinguish Methanol and ethanol.  
(iii) Two organic compounds,  $\text{ROH}$  and  $\text{R}'\text{OH}$ , show different behaviours — one acts as a base while the other acts as an acid. How are the groups  $\text{R}$  and  $\text{R}'$  different from each other?  
(iv) Predict the products of the following reactions :  
a)  $\text{CH}_3 - \text{CH}_2 - \text{CH}_2 - \text{CH}_2 - \text{OH} + \text{SOCl}_2 \rightarrow$   
b)  $\text{CH}_3 - \text{CH}_2 - \text{CH}_2 - \text{O} - \text{CH}_3 + \text{HBr} \rightarrow$

- 32.I. (i) Starch and cellulose both contain glucose units as monomers yet they are structurally different. [ 5 ]  
(ii) The two strands of DNA are complementary to each other.  
(iii) Mention the difference in anomeric forms of glucose and evidence to support the cyclic forms of glucose.  
(iv) How does the formation of glucose pentaacetate during acetylation with acetic anhydride confirm that glucose contains five hydroxyl groups attached to different carbon atoms?  
(v) Which vitamin deficiency causes prolonged bleeding, what is one major source of this vitamin, and what role does dietary fibre play in our body?

(OR)

- 32.II. (i) Which hormone regulates blood sugar in our body, and why should diabetic patients perform mild daily exercises like walking? [ 5 ]
- (ii) A child shows bone deformities, delayed walking, and severe leg pain. Name the nutritional deficiency the child is suffering from.
- (iii) When two nucleotides combine to form a dinucleotide, between which carbon atoms of their pentose sugars is the phosphodiester bond formed?
- (iv) How does the oxidation of glucose with bromine water help to confirm that glucose contains an aldehyde group?
- (v) A person has not eaten for several hours. Enzymes in the liver start breaking down a highly branched carbohydrate stored in the body to release glucose into the blood. Name this stored carbohydrate.

- 33.I.i. The following data were obtained for the reaction: [ 3 ]
- $A + 2B \rightarrow C$


| Experiment | [A]/M | [B]/M | Initial rate of formation of C/M min <sup>-1</sup> |
|------------|-------|-------|--|
| 1          | 0.2   | 0.3   | $4.2 \times 10^{-2}$                               |
| 2          | 0.1   | 0.1   | $6.0 \times 10^{-3}$                               |
| 3          | 0.4   | 0.3   | $1.68 \times 10^{-1}$                              |
| 4          | 0.1   | 0.4   | $2.40 \times 10^{-2}$                              |

- (a) Find the order of reaction with respect to A and B.
- (b) Write the rate law and overall order of reaction.
- (c) Calculate the rate constant (k).
- 33.I.ii. If the activation energy of a reaction is zero. How will the temperature affect the rate of reaction? [ 2 ]

(OR)

33.II.i. Observe the potential energy diagram for the hydrogenation of ethene to give ethane.

[ 3 ]




- (a) For the same process, predict why there is a difference in  $E_{a1}$  and  $E_{a2}$ ?
- (b) Assuming both the reaction are reversible then in which case the backward reaction will be faster?
- (c) Will the  $\Delta G$  value of this reaction be different in the above two paths? Explain.

33.II.ii. Analyse the given graph, drawn between concentration of reactant vs. time.

[ 2 ]

- a) Predict the order of reaction.
- b) Theoretically, can the concentration of the reactant reduce to zero after infinite time? Explain.



**ALL THE BEST**